

Artikel

by Andang Sunarto

Submission date: 02-Sep-2021 03:25PM (UTC+0800)

Submission ID: 1639967751

File name: -sweep_preconditioned_sor_method_for_solving_time-fractional.pdf (1,002.62K)

Word count: 3500

Character count: 17095

3

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/345052188>

6

Application Half-Sweep Preconditioned SOR Method For Solving Time-Fractional Diffusion Equations

Conference Paper · October 2020

CITATIONS

2

READS

53

2 authors:



Andang Sunarto

Universitas Islam Negeri (UIN) Fatmawati Sukarno, Bengkulu, Indonesia

62 PUBLICATIONS 113 CITATIONS

[SEE PROFILE](#)



Jumat Sulaiman

Universiti Malaysia Sabah (UMS)

399 PUBLICATIONS 1,557 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



7

Caputo's finite difference solution of fractional two-point boundary value problems using SOR iteration [View project](#)

9



Baznas [View project](#)

All content following this page was uploaded by Andang Sunarto on 31 October 2020.

The user has requested enhancement of the downloaded file.

6
**Application Half-Sweep Preconditioned SOR Method For
Solving Time-Fractional Diffusion Equations**

Andang Sunarto
IAIN Bengkulu¹⁵ Indonesia
andang99@gmail.com

Jumat Sulaiman
Faculty of Science And Natural Resources
Universiti Malaysia Sabah, Malaysia
jumat@ums.edu.my

Abstract

This research examines the performance of the application Half-Sw³² Preconditioned SOR (HSPSOR) method together with an unconditionally implicit Caputo's time-fractional finite difference approximation equation for solving time-fractional partial diffusion equations (TFPDE's). To do it, the implicit Caputo's time-fractional approximation equations and preconditioned matrix are used to construct the corresponding preconditioned linear system. In addition to that, formulation and application the HSPSOR method are also presented. Based on numerical results of the proposed iterative method, it can be concluded that the proposed iterative method is superior to the Full-Sweep PSOR iterative methods.

Keywords : HSPSOR, Implicit Finite Different, Caputo's, Time-Fractional

1. Introduction

Based on previous studies in (Dey, 1999; Diethelm & Freed, 1999; Gorenflo & Mainardi, 1997; Sene & Fall, 2019; Tam, Wei, & Jin, 2005) many model in mathematical use fractional partial differential equations (FPDEs) to solve fractional problems such as time-fractional partial diffusion equations (TFPDE's). Following to that, there a³¹ several methods used to solve these models. For instance, we have transform method (Sene & Fall, 2019), which is used to obtain analytical and/or numerical solutions of the fractional diffusion equations (FDE's). Other than this method, other researchers have proposed finite difference methods such as explicit and implicit (Chaves, 1998; Evans & Haghight, 1984; Kepczynska, 2005). Also it is pointed out that the explicit iterative methods are conditionally stable. Therefore, to solve the problems of the TFPDE's needs to be discretized. By using the implicit finite difference scheme and Caputo fractional⁷ operator, a linear system at each time level can be construct through the approximation equations. Therefore, Due to the matrix properties of the linear system, iterative methods are the alternative option for efficient solutions

As far as iterative methods are concerned, it can be observed that many researchers such as (Cheng, Huang, & Cheng, 2006; Hackbusch, 2016; Young, 1971) then (Saad, 2003) have proposed and discussed several families of iterative methods. Among the existing proposed methods, the preconditioned iterative methods (Gur¹wardena, Jain, & Snyder, 1991; Salkuyeh & Shamsi, 2012; Shen, Zong, & Shao, 2009) have been widely accepted to be one of the efficient methods for solving linear systems.

Because of the advantages of these iterative methods, the aim of this study is to develop and application the performance of the HSPSOR method to solve (TFPDE's) based on the implicit Caputo's time-fractional finite difference approximation equation. To application the performance of the HSPSOR, we also applying Full-Sweep Preconditioned SOR (FSPSOR).

To demonstrate the performance of HSPSOR, let TFPDE's be given as

$$\frac{\partial^\alpha Z(y,r)}{\partial^\alpha} = a(y) \frac{\partial^2 Z(y,r)}{\partial y^2} + b(y) \frac{\partial Z(y,r)}{\partial y} + c(y) Z(y,r) \quad (1)$$

where $a(y)$, $b(y)$ and $c(y)$ are clear functions or fixed, whereas the parameter α refers to the fractional order TFPDE's derivative.

2. Preliminaries

Previous to construct the discrete equation of Eq (1), in this section given some definitions can be applied for fractional derivative theory

Definition 1. (Sunarto, 2014; Young, 1971) The Riemann-Liouville operator is defined as

$$J^\alpha f(y) = \frac{1}{\Gamma(\alpha)} \int_0^x (y-r)^{\alpha-1} f(r) dr, \alpha > 0, y > 0 \quad (2)$$

Definition 2. (Sunarto, 2014; Young, 1971) The Caputo's operator is defined as

$$D^\alpha f(y) = \frac{1}{\Gamma(m-\alpha)} \int_0^x \frac{f^{(m)}(r)}{(y-r)^{\alpha-m+1}} dr, \alpha > 0 \quad (3)$$

with $m-1 < \alpha \leq m$, $m \in \mathbb{N}$, $y > 0$

To obtain the numerical solution of Eq. (1) with Dirichlet boundary conditions, we get numerical approximations by using the Caputo's derivative definition and consider the non-local fractional TFPDE's derivative operator. This approximation equation can be categorized as unconditionally stable scheme. On strength of Problem (1), the solution domain of the problem has been restricted to the finite space domain $0 \leq y \leq \gamma$ and $0 < \alpha < 1$, whereas α refers to the fractional order of TFPDE's derivative.

In addition to that, consider Initial and boundary (conditions) of Eq (1) be given as

$$Z(x,0) = f(y), \text{ and } Z(0,r) = g_0(r), \quad Z(l,r) = g_1(r), \text{ with } g_0(r), g_1(r), \text{ and } f(y),$$

are given clear functions.

A discretize approximation to TFPDE's Eq. (1) by using Caputo's order- α , is given as [9,11]

$$\frac{\partial^\alpha Z(x,r)}{\partial r^\alpha} = \frac{1}{\Gamma(n-1)} \int_0^\infty \frac{\partial u(y-s)}{\partial r} (r-s)^{-\alpha} ds, \quad r > 0, \quad 0 < \alpha < 1 \quad (4)$$

3. Approximation For Fractional Diffusion Equations

Before constructing the Half-Sweep Caputo's implicit approximation equation, let the Caputo's fractional partial derivative in Eq.(4) be rewritten as

$$D_t^\alpha U_{i,n} \cong \sigma_{\alpha,k} \sum_{j=1}^n \omega_j^{(\alpha)} (U_{i,n-j+1} - U_{i,n-j}) \quad (5)$$

and we have the following expressions

$$\sigma_{\alpha,k} = \frac{1}{\Gamma(1-\alpha)(1-\alpha)k^\alpha}$$

and

$$\omega_j^{(\alpha)} = j^{1-\alpha} - (j-1)^{1-\alpha}.$$

To facilitate us in discretizing the fractional Eq. (1) via the implicit finite difference discretization scheme, Firstly, its the solution domain of the fractional problem is divided uniformly into several subintervals. To do this, we consider several positive integers m and n in which the sizes of subinterval over space and time directions are stated

as $h = \Delta x = \frac{\gamma - 0}{m}$ and $k = \Delta t = \frac{T}{n}$ respectively. According to these sizes of subinterval, we develop the uniformly finite grid network in the solution domain where the grid points over the space interval $[0, \gamma]$ and time interval $[0, T]$ are denoted $x_i = ih$, $i = 0, 1, 2, \dots, m$ and $t_j = jk$, $j = 0, 1, 2, \dots, n$ respectively. Then approximate values of the function $U(x, t)$ at the grid points are labeled as $U_{i,j} = U(x_i, t_j)$. By considering Eq. (5) and the Half-Sweep implicit finite difference discretization scheme, the Half-Sweep Caputo's implicit approximation equation of fractional Eq (1) to the reference grid point at $(x_i, t_j) = (ih, jk)$ is can be shown as

$$\sigma_{\alpha,k} \sum_{j=1}^n \omega_j^{(\alpha)} (U_{i,n-j+1} - U_{i,n-j}) = a_i \frac{1}{4h^2} (U_{i-2,n} - 2U_{i,n} + U_{i+2,n}) + b_i \frac{1}{4h} (U_{i+2,n} - U_{i-2,n}) + c_i U_{i,n}, \quad (6)$$

for $i=2, 4, \dots, m-2$. Actually, the approximation equation (6) also categories as one of family of fully implicit finite difference approximation equations. Also this equation is first order accuracy in time direction and second order in space direction. For simplicity, consider the Half-Sweep approximation equation (6) be rewritten at the specified time level. For instance, we have for $n \geq 2$:

$$\sigma_{\alpha,k} \sum_{j=1}^n \omega_j^{(\alpha)} (U_{i,n-j+1} - U_{i,n-j}) = p_i U_{i-2,n} + q_i U_{i,n} + r_i U_{i+2,n}, \quad (7)$$

where

$$p_i = \frac{a_i}{4h^2} - \frac{b_i}{4h}, \quad q_i = c_i - \frac{a_i}{2h^2}, \quad r_i = \frac{a_i}{4h^2} + \frac{b_i}{4h}.$$

Also, we get for $n = 1$,

$$-p_i U_{i-2,1} + q_i^* U_{i,1} - r_i U_{i+2,1} = f_{i,1}, \quad i = 2, 4, \dots, m-2 \quad (8)$$

where

$$\omega_j^{(\alpha)} = 1, \quad q_i^* = \sigma_{\alpha,k} - q_i, \quad f_{i,1} = \sigma_{\alpha,k} U_{i,1}.$$

Referring to Eq. (8), we can get the tridiagonal linear system that can be constructed in matrix form as

$$AU = f \quad (9)$$

where

$$A = \begin{bmatrix} q_2^* & -r_2 & & & & & \\ -p_4 & q_4^* & -r_4 & & & & \\ & -p_6 & q_6^* & -r_6 & & & \\ & & \circ & \circ & \circ & & \\ & & & -p_{m-2} & q_{m-2}^* & -r_m & \\ & & & & -p_{m-2} & q_{m-2}^* & \left(\left(\frac{m}{2} \right) - 1 \right) \left(\left(\frac{m}{2} \right) - 1 \right) \end{bmatrix},$$

$$U = [U_{2,1} \quad U_{4,1} \quad U_{6,1} \quad \dots \quad U_{m-4,1} \quad U_{m-2,1}]^T, \quad f = [U_{2,1} + p_1 U_{0,1} \quad U_{4,1} \quad U_{6,1} \quad \dots \quad U_{m-4,1} \quad U_{m-2,1} + p_{m-2} U_{m,1}]^T$$

4. Analysis Of Stability

In this section, we have considered the stability analysis of the implicit finite difference approximation equation in Eq.(1). For stability analysis, we will use Von-Neumann's theorem (Langlands & Henry, 2005) and the Lax equivalence theorem (L., Richtmyer, & Morton, 1968). It follows that the numerical solution of the approximation equation in Eq.(1) converges to the exact solution as $h, k \rightarrow 0$.

Theorem 4.1.

4
 The fully implicit numerical method Eq.(1), the solution to Eq.(1) with $0 < \alpha < 1$ on the finite domain $0 \leq x \leq 1$, with zero boundary condition $U(0,t) = U(1,t) = 0$ for all $t \geq 0$, is consistent and unconditionally stable.

Proof. To examine the stability of the proposed method, we find for solution of the form $U_j^n = \xi_n e^{i\omega jh}$, $i = \sqrt{-1}$, ω real. Therefore Eq. (1) becomes

$$\sigma_{\alpha,k} \xi_{n-1} e^{i\omega jh} - \sigma_{\alpha,k} \sum_{j=2}^n \omega_j^{(\alpha)} (\xi_{n-j+1} e^{i\omega jh} - \xi_{n-j} e^{i\omega jh}) = -p_i \xi_n e^{i\omega(j-4)h} + (\sigma_{\alpha,k} - q_i) \xi_n e^{i\omega jh} - r_i \xi_n e^{i\omega(j+4)h} \quad (10)$$

by simplifying and reordering over Eq.(10), we have :

$$\sigma_{\alpha,k} \xi_{n-1} - \sigma_{\alpha,k} \sum_{j=2}^n \omega_j^{(\alpha)} (\xi_{n-j+1} - \xi_{n-j}) = \xi_n ((-p_i - r_i) \cos(\omega h)) + (\sigma_{\alpha,k} - q_i)$$

this can be reduced to :

$$\xi_n = \frac{\xi_{n-1} + \sum_{j=2}^n \omega_j^{(\alpha)} (\xi_{n-j} - \xi_{n-j+1})}{1 + \frac{(p_i + r_i) \cos(\omega h) + q_i}{\sigma_{\alpha,k}}} \quad (11)$$

From Eq.(11), it can be observed that the conducted as

$$\left(1 + \frac{(-p_i - r_i) \cos(\omega h) - q_i}{\sigma_{\alpha,k}} \right) \geq 1, \text{ for all } \alpha, n, \omega, h \text{ and } k \text{ we have } \xi_1 \leq \xi_0. \quad (12)$$

and 33

$$\xi_n \leq \xi_{n-1} + \sum_{j=2}^n \omega_j^{(\alpha)} (\xi_{n-j} - \xi_{n-j+1}), n \geq 2. \quad (13)$$

Thus, for $n=2$, the last inequality implies

$$\xi_2 \leq \xi_1 + \omega_2^{(\alpha)} (\xi_0 - \xi_1)$$

Again repeating the above process, we can get

$$\xi_j \leq \xi_{j-1}, j=1, 2, \dots, n-1.$$

From Eq.(13), we finally have

$$\xi_n \leq \xi_{n-1} + \sum_{j=2}^n \omega_j^{(\alpha)} (\xi_{n-j} - \xi_{n-j+1}) \leq \xi_{n-1}.$$

Since each term in the summation is negative, it shows that the inequalities Eq.(12) and Eq.(13) imply

$$\xi_n \leq \xi_{n-1} \leq \xi_{n-2} \leq \dots \leq \xi_1 \leq \xi_0.$$

Thus,

$$\xi_n = |U_j^n| \leq \xi_0 = |U_j^0| = |f_j|, \text{ which entails } \|U_j^n\| \leq \|f_j\|, \text{ and we have stability.}$$

5. Half-Sweep Preconditioned SOR Method

In relation to the linear system in Equation (8), it is clear that the characteristics of its coefficient matrix are large scale and sparse. Basically in first Section, many scientist have discussed various iteration methods such as (Gunawardena et al., 1991; Hackbusch, 2016; Saad, 2003; Salkuyeh & Shamsi, 2012; Young, 1971). To get numerical solving of the tridiagonal linear system (8), we proposed the Half-Sweep Preconditioned Successive Over Relaxation (HSPSOR) iteration method (Shen et al., 2009; Sunarto, Sulaiman, & Saudi, 2015), to solve any linear systems.

Before applying the HSPSOR iteration method, we need to transform the original linear system (9) into the preconditioned linear system

$$\tilde{A}^* \tilde{Z} = \tilde{f}^* \quad (14)$$

where, $\tilde{A}^* = PAP^T$, $\tilde{f}^* = Pf$, $\tilde{Z} = P^T y$.

Actually, the matrix P is called a preconditioned matrix and defined as (Gunawardena et al., 1991; Sunarto et al., 2015)

$$P = I + S \quad (15)$$

where

$$S = \begin{bmatrix} 0 & -r_1 & 0 & 0 & 0 & 0 \\ 10 & 0 & -r_2 & 0 & 0 & 0 \\ 0 & 0 & 0 & -r_3 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -r_{m-1} \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}_{(m-1) \times (m-1)}$$

and the matrix I is an identical matrix. To develop Half-Sweep PSOR iteration method, let consider the coefficient matrix \tilde{A}^* in equation (9) can be defined as summation of the matrices

$$\tilde{A}^* = Di + Lo + Va \quad (16)$$

with Di , Lo and Va are diagonal, lower triangular and upper triangular matrices. By using Eq.(14) and (16), the formulation of Half-Sweep PSOR iteration method can be rewritten as (Shen et al., 2009; Sunarto et al., 2015)

$$\tilde{y}^{(k+1)} = (D - \omega L)^{-1} [(1 - \omega)D + V\omega] \tilde{y}^{(k)} + (D - \omega L)^{-1} \tilde{f}^* \quad (17)$$

where $\tilde{y}^{(k+1)}$ represents an unknown vector at $(k+1)^{th}$ iteration. The application of the HSPSOR iteration method can be explained in this below.

HSPSOR method

i. First $\tilde{U} \leftarrow 8$ and $\varepsilon \leftarrow 10^{-10}$.

ii. Then $j = 1, 2, K, n$ Implement

a. with $i = 1, 2, K, m-1$ calculate

$$\tilde{y}^{(k+1)} = (D - \omega L)^{-1} [(1 - \omega)D + V\omega] \tilde{y}^{(k)} + (D - \omega L)^{-1} \tilde{f}^* \\ \tilde{Z}^{(k+1)} = P^T \tilde{y}^{(k+1)}$$

b. Next to the convergence test. If the convergence criterion i.e. $\left\| \tilde{y}^{(k+1)} - \tilde{y}^{(k)} \right\| \leq \varepsilon = 10^{-10}$ is satisfied,

go to Step (iii). If not go back to Step (a).

iii Display approximate equation solutions.

6.Evolution Of Numerical Problems

By using approximation (7), we consider one problem of the TFPDE's to test the performance of the Full-Sweep Preconditioned Successive Over-Relaxation (FSPSOR) and Half-Sweep Preconditioned Successive Over-Relaxation (HSPSOR) iteration methods. In order to compare the performance of these proposed iteration methods, three criteria have been considered such as K (number of iterations), Time (in seconds) and Max Absolute Error. three values, where value of $\alpha = 0.25$, $\alpha = 0.50$ and $\alpha = 0.75$. For application of three iterative schemes, the convergence test considered the tolerance error as $\varepsilon = 10^{-10}$.

Let us examine the TFPDE's initial boundary conditions value problem (Ali, S E, Ozgur & Korkmaz, 2013)

$$\frac{\partial^\alpha Z(y, r)}{\partial r^\alpha} = \frac{\partial^2 Z(y, r)}{\partial y^2}, \quad 0 < \alpha \leq 1, 0 \leq y \leq \gamma, r > 0, \quad (18)$$

with the boundary value conditions are $Z(0, r) = \frac{2kr^\alpha}{\Gamma(\alpha+1)}$, $Z(1, r) = 1 + \frac{2kr^\alpha}{\Gamma(\alpha+1)}$, (19)

30
 and the initial value condition $Z(y,0) = y^2$. (20)

Overall results of evolution of numerical problem for equation (18), obtained from application of FSPSOR and HSPSOR iteration methods are recorded in Table 1, where value of mesh sizes, $m = 128, 256, 512, 1024$, and 2048.

7. Conclusions

To In order to get the numerical solution of the TFPDE's problems equation, this study give the derivation of the implicit Caputo's finite difference approximation equations in which this approximation equation leads a tridiagonal linear system. Via all experimental results by imposing the FSPSOR and HSPSOR iteration methods, it is clear at $\alpha = 0.25$ that K (number of iterations) have declined approximately by 64.27-96.14% conforms to the HSPSOR iteration method compared with FSPSOR methods. Then for Time, application of HSPSOR iteration method are much faster about 25.84-94.18% than the FSPSOR methods. It can be also observed in Table 1 that the HSPSOR method requires the least amount for K (number of iterations) and Time $\alpha = 0.25$ as compared with FSPSOR iteration methods. According to the accuracy of both iteration methods, it can be concluded that their numerical solutions of TFPDE's are in good agreement.

References

- Ali, S E, Ozgur, B., & Korkmaz, E. (2013). Analysis Of Fractional Partial Differential Equations By Taylor Series Expansion. *Boundary Value Problem, A Springer Open*, 68.
- Chaves, A. S. (1998). A fractional diffusion equation to describe Lévy flights. *Physics Letters A*, 239(1-2), 13-16. [https://doi.org/10.1016/s0375-9601\(97\)00947-x](https://doi.org/10.1016/s0375-9601(97)00947-x)
- Cheng, G. H., Huang, T. Z., & Cheng, X. Y. (2006). Preconditioned Gauss-Seidel type iterative method for solving linear systems. *Applied Mathematics and Mechanics (English Edition)*, 27(9), 1275-1279. <https://doi.org/10.1007/s10483-006-0915-1>
- Dey, S. K. (1999). A novel explicit finite difference scheme for partial differential equations. *Mathematical Modelling and Analysis*, 4(1), 70-78. <https://doi.org/10.1080/13926292.1999.9637112>
- Diethelm, K., & Freed, A. D. (1999). On the Solution of Nonlinear Fractional-Order Differential Equations Used in the Modeling of Viscoplasticity. In *Scientific Computing in Chemical Engineering II* (pp. 217-224). https://doi.org/10.1007/978-3-642-60185-9_24
- Evans, D. J., & Haggihl, R. S. (1984). Explicit Group Versus Implicit Line Iterative Methods. *International Journal of Computer Mathematics*, 16(4), 261-316. <https://doi.org/10.1080/00207168408803442>
- Gorenflo, R., & Mainardi, F. (1997). Fractional Calculus. In *Fractals and Fractional Calculus in Continuum Mechanics* (pp. 223-276). https://doi.org/10.1007/978-3-7091-2664-6_5
- Gunawardena, A. D., Jain, S. K., & Snyder, L. (1991). Modified iterative methods for consistent linear systems. *Linear Algebra and Its Applications*, 154-156(C), 123-143. [https://doi.org/10.1016/0024-3795\(91\)90376-8](https://doi.org/10.1016/0024-3795(91)90376-8)
- Hackbusch, W. (2016). Iterative solution of large sparse systems of equations. In *Applied Mathematical Sciences (Switzerland)* (Vol. 95). <https://doi.org/10.2307/2153387>
- I., E., Richtmyer, R. D., & Morton, K. W. (1968). Difference Methods for Initial-Value Problems. *Mathematics of Computation*, 22(102), 465. <https://doi.org/10.2307/2004698>
- Kepczynska, A. (2005). Implicit difference methods for first-order partial differential functional equations. *Nonlinear Oscillations*, 8(2), 198-213. <https://doi.org/10.1007/s11072-005-0049-z>
- Langlands, T. A. M., & Henry, B. I. (2005). The accuracy and stability of an implicit solution method for the fractional diffusion equation. *Journal of Computational Physics*, 205(2), 719-736. <https://doi.org/10.1016/j.jcp.2004.11.025>
- Saad, Y. (2003). 9. Preconditioned Iterations. In *Iterative Methods for Sparse Linear Systems* (pp. 261-281). <https://doi.org/10.1137/1.9780898718003.ch9>
- Salkuyeh, D. K., & Shamsi, S. (2012). A preconditioner for the SOR-like method for the augmented systems. *Chiang Mai Journal of Science*, 39(2), 213-221.
- Sene, N., & Fall, A. N. (2019). Homotopy Perturbation p-Laplace Transform Method and Its Application to the Fractional Diffusion Equation and the Fractional Diffusion-Reaction Equation. *Fractal and Fractional*, 3(2), 14. <https://doi.org/10.3390/fractalfract3020014>
- Shen, H. L., Zong, Y., & Shao, X. H. (2009). Preconditioned SOR-type iterative methods for solving linear systems. *Dongbei Daxue Xuebao/Journal of Northeastern University*, 30(8), 1213-1216.
- Sunarto, A. (2014). Borneo science 34: march 2014. *Borneo Science*, (March), 34-42.

- Sunarto, A., Sulaiman, J., & Saudi, A. (2015). Solving the time fractional diffusion equations by the Half-Sweep SOR iterative method. In *Proceedings - 2014 International Conference on Advanced Informatics: Concept, Theory and Application, ICAICTA 2014*. <https://doi.org/10.1109/ICAICTA.2014.7005953>
- Tam, H. S., Wei, Y. M., & Jin, X. Q. (2005). A note on preconditioning for M-matrix. *Applied Mathematics Letters*, 18(10), 1137–1142. <https://doi.org/10.1016/j.aml.2005.02.001>
- Young, D. . (1971). *Iterative solution of large linear systems*.

TABLE 1. Comparison between number of iterations (K), the computational time (seconds) and maximum absolute errors (MAE) for both iterative methods at $\alpha = 0.25, 0.50, 0.75$

M	Method	$\alpha = 0.25$			$\alpha = 0.50$			$\alpha = 0.75$		
		K	Time	MAE	K	Time	MAE	K	Time	MAE
128	FSPSOR	281	2.24	9.95e-05	229	1.95	9.84e-05	164	1.63	1.29e-04
	HSPSOR	156	1.21	9.95e-05	137	1.06	9.84e-05	132	0.99	1.30e-04
256	FSPSOR	1428	16.9	9.95e-05	1171	12.61	9.84e-05	814	8.90	1.29e-04
	HSPSOR	778	4.89	9.95e-05	486	3.31	9.84e-05	211	1.90	1.30e-04
512	FSPSOR	5524	113.86	9.96e-05	4520	91.37	9.84e-05	3137	61.98	1.30e-04
	HSPSOR	3006	32.07	9.96e-05	1929	20.66	9.84e-05	931	10.60	1.30e-04
1024	FSPSOR	20574	817.59	9.98e-05	16842	662.23	9.87e-05	11695	456.23	1.30e-04
	HSPSOR	11216	236.40	9.98e-05	7245	155.08	9.87e-05	3551	74.13	1.30e-04
2048	FSPSOR	75580	3043.59	1.01e-04	61941	2894.70	9.90e-05	43070	977.10	1.30e-04
	HSPSOR	40270	1835.90	9.98e-05	25487	1320.76	9.84e-05	13208	552.95	1.30e-04

Biography / Biographies

Andang Sunarto is an Lecturer, Institut Agama Islam Negeri, Bengkulu, Indonesia

Jumat Sulaiman is a Lecturer in Faculty of Mathematics And Natural Resources, Universiti Malaysia Sabah, Malayisa

Artikel

ORIGINALITY REPORT

21 %
SIMILARITY INDEX

14 %
INTERNET SOURCES

14 %
PUBLICATIONS

8 %
STUDENT PAPERS

PRIMARY SOURCES

- | | | |
|---|--|-----|
| 1 | M. S. Muthuvalu, J. Sulaiman. "Numerical solutions of second kind Fredholm integral equations using Half-Sweep Geometric Mean method", 2008 International Symposium on Information Technology, 2008
Publication | 2 % |
| 2 | Submitted to Universiti Malaysia Perlis
Student Paper | 2 % |
| 3 | es.scribd.com
Internet Source | 1 % |
| 4 | Submitted to Universiti Sains Malaysia
Student Paper | 1 % |
| 5 | ejpam.com
Internet Source | 1 % |
| 6 | ieomsociety.org
Internet Source | 1 % |
| 7 | www.ijettjournal.org
Internet Source | 1 % |

8	A Sunarto, J Sulaiman, J V L Chew. "Performance of FSPAOR iteration for solving one-dimensional space-fractional diffusion equation", Journal of Physics: Conference Series, 2021 Publication	1 %
9	dokumen.pub Internet Source	1 %
10	www.elpa.it Internet Source	1 %
11	Submitted to Universitas Islam Indonesia Student Paper	1 %
12	F A Muhiddin, J Sulaiman, A Sunarto. "Grünwald Implicit Solution of One-Dimensional Time-Fractional Parabolic Equations Using HSKSOR Iteration", Journal of Physics: Conference Series, 2020 Publication	1 %
13	www.scribd.com Internet Source	1 %
14	N. A. M. Ali, J. Sulaiman, A. Saudi, N. S. Mohamad. "Chapter 42 The Similarity Finite Difference Solutions for Two-Dimensional Parabolic Partial Differential Equations via SOR Iteration", Springer Science and Business Media LLC, 2021 Publication	1 %

15	Submitted to Universiti Malaysia Sabah Student Paper	1 %
16	benthamopen.com Internet Source	1 %
17	www.astesj.com Internet Source	1 %
18	www.thaiscience.info Internet Source	1 %
19	Langlands, T.A.M.. "The accuracy and stability of an implicit solution method for the fractional diffusion equation", Journal of Computational Physics, 20050520 Publication	<1 %
20	fbe.erciyes.edu.tr Internet Source	<1 %
21	www.slideshare.net Internet Source	<1 %
22	Fatihah Anas Muhiddin, Jumat Sulaiman, Andang Sunarto. "Chapter 24 Solving Time-Fractional Parabolic Equations with the Four Point-HSEGKSOR Iteration", Springer Science and Business Media LLC, 2021 Publication	<1 %
23	Young-II Lim, Stem Bay Jrgensen. "Distributed Dynamic Models and Computational Fluid Dynamics", Wiley, 2006	<1 %

24	rxiv.org Internet Source	<1 %
25	www.koreascience.or.kr Internet Source	<1 %
26	Submitted to University of Bradford Student Paper	<1 %
27	www.gjetr.org Internet Source	<1 %
28	Eng Jeng Hong, Azali Saudi, Jumat Sulaiman. "Application of SOR Iteration for Poisson Image Blending", Proceedings of the International Conference on High Performance Compilation, Computing and Communications - HP3C-2017, 2017 Publication	<1 %
29	core.ac.uk Internet Source	<1 %
30	file.scirp.org Internet Source	<1 %
31	www.scielo.org.mx Internet Source	<1 %
32	www.springerprofessional.de Internet Source	<1 %

33

István Kovács, Aleksander Malnič, Dragan Marušič, Štefko Miklavič. "Transitive group actions: (im)primitivity and semiregular subgroups", Journal of Algebraic Combinatorics, 2014

Publication

<1 %

34

Texts in Computational Science and Engineering, 2003.

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On